

Heavy quark masses from lattice QCD*

Jochen Heitger^a

^aWestfälische Wilhelms-Universität Münster, Institut für Theoretische Physik,
Wilhelm-Klemm-Strasse 9, D-48149 Münster, Germany

I outline the basic strategies for the computation of charm and bottom quark masses by means of lattice QCD, where particular emphasis is placed on the non-perturbative renormalization of the effective theory for the b-quark in heavy-light systems. A few selected results in the quenched approximation are reviewed, and the current status of extending these calculations to QCD with dynamical quarks is summarized.

1. Introduction

The fundamental parameters of QCD are the strong coupling constant and the quark masses. In particular, high-precision tests of the SM critically depend on the masses of the charm and bottom quarks, and the non-perturbative nature of lattice QCD, which allows to directly compute hadronic observables to match experimental inputs, is especially suited to determine them.

Their ab initio lattice calculation faces systematic errors, such as cutoff and finite-volume effects, too large dynamical light quark masses and the omission of some (or, in quenched QCD, even all) sea quark flavours. Among these, the problem of discretization errors proportional to powers of the bare quark mass is of utmost relevance in case of systems containing a heavy quark, when a Wilson-like fermion action is employed. Requiring $am_{q,h} \ll 1$, it is then obvious that simulating a b-quark at its physical mass of ≈ 5 GeV in a space-time volume of $L^4 = (2 \text{ fm})^4$ would demand $\gg 50$ lattice points per direction, an effort beyond the computing resources available today.

In this situation, an attractive possibility is to recourse to an *effective theory* for the b-quark. Extracting physical predictions from it, however, involves a matching to QCD, which becomes a severe source of uncertainty when performed perturbatively, for in the continuum limit results are power divergent. Before coming to the effective theory framework to deal with heavy-light sys-

tems on the lattice (Section 3), let us consider the c-quark, where direct simulations are just doable.

2. The charm quark's mass

Physics in the region around the charm quark, which is by a factor of about 4 lighter than the b-quark, can be studied with conventional lattice QCD methods, provided that — assuming Wilson fermions here — the leading $O(a)$ cutoff effects have been eliminated non-perturbatively and a range of lattice resolutions $a^{-1} \approx (2 - 4) \text{ GeV}$ is covered in the simulations. This is nicely illustrated by the benchmark calculation [1] of the mass of the c-quark in the quenched approximation, which accounts for all systematic errors except for the neglect of dynamical quark. Following the strategy and techniques used to compute m_s [2], and $O(am_{q,c})$ effects $\propto (b_A - b_P)$ being removed [3], the renormalization group invariant (RGI) c-quark mass was calculated on 4 lattices with spacings $a = (0.1 - 0.05) \text{ fm}$. In addition to the definition of the (non-perturbatively) renormalized quark mass via the heavy-light PCAC relation, two further definitions were considered, differing by $O(a^2)$ discretization errors. The agreement of all three definitions in the continuum limit found in [1] thus provides a clear demonstration that on lattices of the chosen size (from $16^3 \times 32$ to $32^3 \times 64$) one can control the discretization errors carefully by an extrapolation.

This is confirmed by including a further gauge field ensemble in a volume of $48^3 \times 96$ with an even finer resolution of $a = 0.03 \text{ fm}$, which was first generated in the context of [4] to investigate,

*Invited talk at the Workshop on e^+e^- Collisions from Φ to Ψ (PHIPSI08), April 7–10, 2008, INFN Frascati, Italy.




Figure 1. Continuum extrapolation of the RGI c -mass for $N_f = 0$, covering $a = (0.1 - 0.03)$ fm; see [5] for details. The length scale $r_0 = 0.5$ fm is derived from the force between static quarks.

within quenched QCD, the size of cutoff effects in the charm sector thoroughly also for other observables such as the D_s -meson decay constant. Figure 1 shows an update of the continuum extrapolation of the RGI charm mass after a significant increase of the statistics compared to [1,4], particularly for the smallest a [5]. The final result of this analysis, $\overline{m}_c^{\overline{\text{MS}}}(\overline{m}_c) = 1268(24)$ MeV, is compatible with the earlier result of ref. [1].

However, from the figure it is also evident that a controlled assessment of the cutoff effects will not be possible for much heavier quarks. Hence, for b -quarks other approaches must be devised, and one will be discussed in the next section.

Charm-quark mass computations in QCD with dynamical flavours are not yet in the stage such that a solid continuum limit extrapolation can be performed [6,7]. Since non-perturbative estimates for all the necessary improvement coefficients and renormalization factors in the two-flavour Wilson theory have become available now [8,9,10,11], $N_f = 2$ calculations of masses and matrix elements in the charm sector along the lines of the quenched studies are being started [12].

For other recent determinations of $\overline{m}_c^{\overline{\text{MS}}}$ using QCD sum rules and current-current correlators in lattice and continuum QCD, see refs. [13,14].

3. Non-perturbative HQET

Heavy Quark Effective Theory (HQET) at zero velocity on the lattice [15] offers a reliable solution to the problem of dealing with the two disparate intrinsic scales encountered in heavy-light systems involving the b -quark, i.e., the lattice spacing a , which has to be much smaller than

$1/m_b$ to allow for a fine enough resolution of the states in question, and the linear extent L of the lattice volume, which has to be large enough for finite-size effects to be under control.

Since the heavy quark mass (m_b) is much larger than the other scales such as its 3-momentum or $\Lambda_{\text{QCD}} \sim 500$ MeV, HQET relies upon a systematic expansion of the QCD action and correlation functions in inverse powers of the heavy quark mass around the static limit ($m_b \rightarrow \infty$). The lattice HQET action, S_{HQET} , at $\mathcal{O}(1/m_b)$ reads:

$$a^4 \sum_x \bar{\psi}_h \{ D_0 + \delta m - \omega_{\text{kin}} \mathbf{D}^2 - \omega_{\text{spin}} \boldsymbol{\sigma} \mathbf{B} \} \psi_h,$$

with ψ_h satisfying $P_+ \psi_h = \psi_h$, $P_+ = \frac{1+\gamma_0}{2}$, and the parameters ω_{kin} and ω_{spin} being formally $\mathcal{O}(1/m_b)$. At leading order (static limit), where the heavy quark acts only as a static colour source and the light quarks are independent of the heavy quark's flavour and spin, the theory is expected to have $\sim 10\%$ precision, while this reduces to $\sim 1\%$ at $\mathcal{O}(1/m_b)$ representing the interactions due to motion and spin of the heavy quark. As crucial advantage (e.g., over NRQCD), HQET treats the $1/m_b$ -corrections to the static theory as space-time insertions in correlations functions, viz.

$$\langle \mathcal{O} \rangle = \langle \mathcal{O} \rangle_{\text{stat}} + a^4 \sum_x \{ \omega_{\text{kin}} \langle \mathcal{O} \mathcal{O}_{\text{kin}}(x) \rangle_{\text{stat}} + \omega_{\text{spin}} \langle \mathcal{O} \mathcal{O}_{\text{spin}}(x) \rangle_{\text{stat}} \}$$

for multi-local fields $\mathcal{O}$, where $\langle \mathcal{O} \rangle_{\text{stat}}$ denotes the expectation value in the static approximation and $\mathcal{O}_{\text{kin}}$ and $\mathcal{O}_{\text{spin}}$ are given by $\bar{\psi}_h \mathbf{D}^2 \psi_h$ and $\bar{\psi}_h \boldsymbol{\sigma} \mathbf{B} \psi_h$. In this way, HQET at a given order is (power-counting) renormalizable and its continuum limit well defined, once the mass counterterm δm and the coefficients ω_{kin} and ω_{spin} are fixed non-perturbatively by a matching to QCD.

Still, for lattice HQET and its numerical applications to lead to precise results with controlled systematic errors in practice, two shortcomings had to be left behind first.

1. The exponential growth of the noise-to-signal ratio in static-light correlation functions, which can be overcome by a clever modification of the Eichten-Hill discretization of the static action [16].

2. As in HQET mixings among operators of different dimensions occur, the power-divergent additive mass renormalization $\delta m \sim g_0^2/a$ already affects its leading order. Unless HQET is renormalized non-perturbatively [17], this divergence — and further ones $\sim g_0^2/a^2$ arising at $\mathcal{O}(1/m_b)$ — imply that the continuum limit does not exist owing to a remainder, which, at any finite perturbative order [18,19], diverges as $a \rightarrow 0$. A general solution to this theoretically serious problem was worked out and implemented for a determination of the b-quark's mass in the static and quenched approximations as a test case in ref. [20]. It is based on a *non-perturbative matching of HQET and QCD in finite volume*.

In Section 3.1 I review the HQET computation of the mass of the b-quark including the $\mathcal{O}(1/m_b)$ terms for $N_f = 0$ [21], while Section 3.2 briefly summarizes the present status of the ongoing project to extend this to the more realistic case of two-flavour QCD [22]. A first application of the strategy of [20] to the computation of the B-meson decay constant can be found in [23].

A promising, not unrelated approach to lattice heavy-light systems derives from the step scaling method in the relativistic theory proposed in [24], where extrapolations in the heavy quark mass of step scaling functions (SSFs) as finite-size effects of proper observables, combined with SSFs calculated in HQET, turn into safer interpolations [25].

3.1. The b-quark mass in HQET at $\mathcal{O}(\frac{1}{m_b})$

As pointed out in [26], let us first note that in order not to spoil the asymptotic convergence of the series, the matching must be done non-perturbatively — at least for the leading, static piece — as soon as the $1/m_b$ -corrections are included, since as $m_b \rightarrow \infty$ the *perturbative* truncation error from the matching coefficient of the static term becomes much larger than the power corrections $\sim \Lambda_{\text{QCD}}/m_b$ of the HQET expansion.

In the framework introduced in ref. [20], matching and renormalization are performed simultaneously *and* non-perturbatively. The general strategy, illustrated in figure 2, can be explained as follows. Starting from a finite volume with $L_1 \approx$




Figure 2. *Idea of lattice HQET computations via a non-perturbative determination of HQET parameters from small-volume QCD simulations. Arrows indicate steps to be repeated at smaller a to reach a continuum limit. (Drawing from [26].)*

0.5 fm, one chooses lattice spacings a sufficiently smaller than $1/m_b$ such that the b-quark propagates correctly up to controllable discretization errors of order a^2 . The relation between the RGI and the bare mass in QCD being known, suitable finite-volume observables $\Phi_k(L_1, M_b)$ can be calculated as a function of the RGI b-quark mass, M_b , and extrapolated to the continuum limit. Next, the power-divergent subtractions are performed non-perturbatively by a set of matching conditions, in which the results obtained for Φ_k are equated to their representation in HQET (r.h.s. of figure 2). At the same physical value of L_1 but for resolutions $L_1/a = \mathcal{O}(10)$, the previously computed heavy-quark mass dependence of $\Phi_k(L_1, M_b)$ in finite-volume QCD may be exploited to determine the bare parameters of HQET for $a \approx (0.025 - 0.05)$ fm. In order to evolve the HQET observables to large volumes, where contact with experiments can be made, one also computes them at these lattice spacings in a larger volume, $L_2 = 2L_1$. The resulting relation between $\Phi_k(L_1)$ and $\Phi_k(L_2)$ is encoded in associated SSFs σ_k , as indicated in figure 2. Finally, the knowledge of $\Phi_k(L_2, M_b)$ and employing resolutions $L_2/a = \mathcal{O}(10)$ fixes the bare parameters of the effective theory for $a \approx (0.05 - 0.1)$ fm so that a connection to lattice spacings is established, where large-volume observables, such as

the B-meson mass or decay constant, can be calculated (l.h.s. of figure 2). This sequence of steps yields an expression of m_B (the physical input) as a function of M_b via the quark mass dependence of $\Phi_k(L_1, M_b)$, which eventually is inverted to arrive at the desired value of the RGI b-mass within HQET. The whole construction is such that the continuum limit can be taken for all pieces.

More specifically, upon restricting to spin-averaged quantities to get rid of the contributions proportional to ω_{spin} , the task is to fix δm and ω_{kin} non-perturbatively by performing a matching to QCD. For sensible definitions of the required matching observables, Φ_1 and Φ_2 , we work with the Schrödinger functional (SF), i.e. QCD with Dirichlet boundary conditions in time and periodic ones in space (up to a phase θ for the fermions): $\Phi_1^{\text{QCD}}(L, m_h)$ exploits the sensitivity of SF correlators to θ and $\Phi_2^{\text{QCD}}(L, m_h) \equiv L\Gamma_1(L, m_h)$, where Γ_1 is a finite-volume effective energy. When expanded in HQET², $\Phi_1^{\text{HQET}}(L)$ is given by ω_{kin} times a quantity defined in the effective theory (called $R_1^{\text{kin}}(L, \theta, \theta')$), whereas $\Phi_2^{\text{HQET}}(L)$ is a function of ω_{kin} and $m_{\text{bare}} = \delta m + m_h$ involving two other HQET quantities, $\Gamma_1^{\text{stat}}(L)$ and $\Gamma_1^{\text{kin}}(L)$. According to the strategy sketched above, by equating $\Phi_k^{\text{QCD}}(L_1, m_h)$ and $\Phi_k^{\text{HQET}}(L_1)$ one can determine the bare parameters m_{bare} and ω_{kin} as functions of m_h at the lattice spacings belonging to the volume L_1^4 . To use the spin-averaged B-meson mass, m_B^{av} , as phenomenological input, the Φ_k are evolved to larger volumes through proper SSFs [21], where the resulting $\Phi_k^{\text{HQET}}(2L_1, m_h)$ still carry the dependence on m_h inherited from the matching to QCD in L_1^4 . After 2 evolution steps (and taking continuum limits), linear extents of $\gtrsim 1.5$ fm are reached, and m_{bare} and ω_{kin} , expressed in terms of SSFs, $\Phi_k^{\text{QCD}}(L_1, m_h)$ as well as R_1^{kin} , Γ_1^{stat} and Γ_1^{kin} , are obtained — again as functions of m_h . Now, the b-quark mass is extracted by solving

$$m_B^{\text{av}} = E^{\text{stat}} + \omega_{\text{kin}}(m_h)E^{\text{kin}} + m_{\text{bare}}(m_h) \quad (1)$$

for m_h , with $E^{\text{stat}} = \lim_{L \rightarrow \infty} \Gamma_1^{\text{stat}}$ and $E^{\text{kin}} = -\langle B | a^3 \sum_{\mathbf{z}} \mathcal{O}_{\text{kin}}(0, \mathbf{z}) | B \rangle_{\text{stat}}$. All quantities enter-

²Here, $\delta m = 0$ in the action; its contribution is accounted for in the overall energy shift m_{bare} in HQET versus QCD.

ing eq. (1) having a continuum limit either in QCD or HQET implies that all power divergences have been subtracted non-perturbatively.




Figure 3. Graphical solution of eq. (2) in the quenched approximation [21]. The quantity used in the finite-volume matching step is $\Phi_2^{\text{QCD}}(L_1, M) = L_1 \Gamma_1(L_1, M)$, and $z \equiv L_1 M$.

In case of the leading-order, static approximation, where only m_{bare} needs to be determined, the small- and large-volume matching conditions simplify to $\Gamma_1(L_1, m_h) = \Gamma_1^{\text{stat}}(L_1) + m_{\text{bare}}$ and $m_B^{\text{av}} = E^{\text{stat}} + m_{\text{bare}}$, respectively. To be able to replace m_{bare} in the latter by the former, we bridge the volume gap in two steps by inserting a SSF $\sigma_m(L_1) = 2L_1[\Gamma_1^{\text{stat}}(2L_1) - \Gamma_1^{\text{stat}}(L_1)]$ and arrive at the *master equation*

$$L_1 [m_B^{\text{av}} - (E^{\text{stat}} - \Gamma_1^{\text{stat}})] - \frac{\sigma_m(L_1)}{2} = L_1 \Gamma_1, \quad (2)$$

where $\Gamma_1 = \Gamma_1(L_1, m_h)$ stems from L_1^4 -QCD and any reference to bare parameters has finally disappeared. Its graphical solution is reproduced in figure 3 and yields $M_b^{\text{stat}} = 6.806(79)$ GeV.

For the details on the (technically more involved) inclusion of the sub-leading $1/m_b$ -effects, which exploits the freedom in choices for the angle θ as well as an alternative set of matching observables, I refer to ref. [21]. Here, I just quote their final result $\overline{m}_b^{\text{MS}}(\overline{m}_b) = 4.347(48)$ GeV with the remark that, upon including the $1/m_b$ -terms, differences among the static results w.r.t. the matching condition chosen are gone, which signals practically negligible higher-order corrections.

3.2. Status in two-flavour QCD

The renormalization of HQET through the non-perturbative matching to $N_f = 2$ QCD in fi-

nite volume, to perform the power-divergent subtractions, is under way [22]. As an important prerequisite, the non-perturbative relation between the RGI and subtracted bare heavy quark mass calculated in [10,11] enables to fix RGI heavy quark masses in the matching volume L_1^4 :

$$L_1 M = Z_M(g_0) Z(g_0) (1 + b_m(g_0) a m_{q,h}) L_1 m_{q,h}.$$

The extent L_1 is defined via a constant SF coupling, $\bar{g}^2(L_1/2) = 2.989$, and the PCAC masses of the dynamical light quarks are tuned to zero.

Figure 4 shows an example for the mass dependence of a finite-volume QCD observable in the continuum limit, which enters the matching step.




Figure 4. Preliminary continuum limit (left) and z -dependence (right) of the spin-averaged B -meson energy in finite-volume QCD for $N_f = 2$.

4. Outlook

Thanks to technical and conceptual advances, most notably regarding non-perturbative $O(a)$ improvement and renormalization in QCD and HQET, heavy-light physics has recently seen a significant progress. Since most current simulations control other systematics such as dynamical quark effects and the continuum extrapolation, more precise results for the c - and b -quark masses can be expected within the next years.

Acknowledgments. I am indebted to my colleagues B. Blossier, P. Fritzsche, N. Garron, G. von Hippel, H. Meyer, M. Della Morte, S. Schäfer, H. Simma and R. Sommer from the ALPHA Collaboration and to G. De Divitiis, A. Jüttner and N. Tantalo for an enjoyable, fruitful collaboration. We thank NIC/DESY for allocating computer time on the APE and BlueGene/L&P computers to this project. Support by the European Community through EU Contract No. MRTN-CT-2006-035482, “FLAVIANet” and by the DFG under grant HE 4517/2-1 is acknowledged.

REFERENCES

1. J. Rolf and S. Sint, JHEP 12 (2002) 007, hep-ph/0209255.
2. J. Garden et al., Nucl. Phys. B571 (2000) 237, hep-lat/9906013.
3. M. Guagnelli et al., Nucl. Phys. B595 (2001) 44, hep-lat/0009021.
4. A. Jüttner, hep-lat/0503040.
5. J. Heitger and A. Jüttner, arXiv:0812.2200.
6. A. Dougall, C.M. Maynard and C. McNeile, JHEP 01 (2006) 171, hep-lat/0508033.
7. B. Blossier, G. Herdoiza and S. Simula, PoS LAT2007 (2007) 346, arXiv:0710.1414.
8. M. Della Morte, R. Hoffmann and R. Sommer, JHEP 03 (2005) 029, hep-lat/0503003.
9. M. Della Morte et al., JHEP 07 (2005) 007, hep-lat/0505026; arXiv:0807.1120.
10. M. Della Morte et al., Nucl. Phys. B729 (2005) 117, hep-lat/0507035.
11. M. Della Morte et al., arXiv:0710.1188; G.M. de Divitiis et al., in preparation.
12. J. Heitger et al., arXiv:0810.0214.
13. J. Kühn, M. Steinhauser and C. Sturm, Nucl. Phys. B778 (2007) 192, hep-ph/0702103.
14. I. Allison et al., (2008), arXiv:0805.2999.
15. E. Eichten and B. Hill, Phys. Lett. B234 (1990) 511.
16. M. Della Morte, A. Shindler and R. Sommer, JHEP 08 (2005) 051, hep-lat/0506008.
17. L. Maiani, G. Martinelli and C.T. Sachrajda, Nucl. Phys. B368 (1992) 281.
18. G. Martinelli and C.T. Sachrajda, Nucl. Phys. B559 (1999) 429, hep-lat/9812001.
19. F.D. Renzo and L. Scorzato, JHEP 02 (2001) 020, hep-lat/0012011.
20. J. Heitger and R. Sommer, JHEP 02 (2004) 022, hep-lat/0310035.
21. M. Della Morte et al., JHEP 01 (2007) 007, hep-ph/0609294.
22. M. Della Morte et al., arXiv:0810.3166.
23. B. Blossier et al., PoS LAT2007 (2007) 245, arXiv:0710.1553.
24. G.M. de Divitiis et al., Nucl. Phys. B675 (2003) 309, hep-lat/0305018.
25. D. Guazzini, R. Sommer and N. Tantalo, JHEP 01 (2008) 076, arXiv:0710.2229.
26. R. Sommer, hep-lat/0611020.